


Quick Review

$A \cup B$, union

consists of
all elements in
 A or in B


$$U \approx V$$



$A - B$, difference

$$A - B$$


consists of
all elements in
 A but not in
 B



$A \cap B$, intersection


consists of
all elements
in A and B

$$\cap \approx \wedge$$



\bar{A} , complement

First specify the
universe U ,
the set that
consists of everyth-
ing in U not in A .



$$f: A \rightarrow B$$




domain



codomain
≠ range


injectivity:

0



surjectivity:


①



bijection


$$- f: A \rightarrow B$$

$$|A| = |B|$$



the image of a set $A \subseteq X$ is the set $f(A) = \{ f(a) \mid a \in A \}$

the pre-image of a set $B \subseteq Y$ is




$$f^{-1}(B) = \{ a \mid a \in X, f(a) \in B \}$$

$f(x) = \frac{1}{x}$ a function?

$f: \mathbb{R} \rightarrow \mathbb{R}$ then no.

$f: \mathbb{R} \setminus \{0\} \rightarrow \mathbb{R}$ then yes.



$f(2)$ can't
be two
things.

X domain

Y codomain

$\forall x \in X, f(x)$ exists
and is
unique.

Discussion 1D Problem 1

(a) If $A = \{1, 2, 3, 4\}$, what is $\mathcal{P}(A)$?

The set of all subsets of $\{1, 2, 3, 4\}$, which is tedious to write out.

(b) If B is a set, describe $\mathcal{P}(B)$ using set comprehension.

$$\{b \mid b \subseteq B\}.$$

(c) What is $\mathbb{R} \cap \mathcal{P}(A)$?

\emptyset , as the elements of \mathbb{R} are numbers and the elements of $\mathcal{P}(A)$ are sets

(d) What is $\mathbb{R} \cap \mathbb{Z}$?

\mathbb{Z} , since the integers are a subset of the reals.

(e) What is in $\mathbb{N} \cup \mathbb{Q}$?

\mathbb{Q} , since the naturals are a subset of the rationals.

(f) What kind of numbers are in $\mathbb{R} \setminus \mathbb{Q}$?

Irrational numbers.

(g) If $S \subseteq T$, what is $S \setminus T$.

\emptyset , as T contains all of S .

Discussion 1 D Problem 2

Let X, Y be sets, and $f: X \rightarrow Y$ be a function.

- If $A \subseteq X$, define the image of A under f as the set



$$f(A) := \{ f(x) \mid x \in A \}.$$

- If $B \subseteq Y$, define the pre-image of B under f as the set

$$f^{-1}(B) := \{ x \mid x \in X \wedge f(x) \in B \}.$$

- (a) If $B \subseteq Y$, prove that $f(f^{-1}(B)) \subseteq B$.

Idea: Prove that ~~$\forall b \in B$,~~
 $\forall b \in \underline{f(f^{-1}(B))}$, $b \in B$



Idea: Prove $\forall b \in f(f^{-1}(B)), b \in B$

Let $b \in f(f^{-1}(B))$ be arbitrary.

Then ~~$f(\exists a \in f^{-1}(B)$~~ such that
 $f(a) = b$. Since $a \in f^{-1}(B)$, $f(a)$
 $\in B$. $b = f(a) \in B$.

(b/c) Let $A \subseteq X$. Prove that $A \subseteq f^{-1}(f(A))$ and provide an example where $A \neq f^{-1}(f(A))$


Idea: $\forall a \in A$, prove that $a \in f^{-1}(f(A))$

Since $a \in A$, it follows that $f(a) \in f(A)$. By the definition of the preimage, $a \in f^{-1}(f(A))$ because $f(a) \in f(A)$.

$$f(x) = x^2, f: \mathbb{R} \rightarrow \mathbb{R}.$$

$$A = [0, 1].$$

$$f(A) = [0, 1]$$



$$f^{-1}(f(A))$$

Discussion 1 D Problem 3

Consider the function

$$f(x) = \begin{cases} x & \text{if } x \geq 1 \\ x^2 & \text{if } -1 \leq x < 1 \\ 2x + 3 & \text{otherwise.} \end{cases}$$

(a) If the domain / codomain of f are \mathbb{N} ,
is f

- Injective?

Yes, as on \mathbb{N} , $f(x) = x$, so if
 $f(a) = f(b)$, then $a = b$.

- Surjective?

Yes. If $n \in \mathbb{N}$ is arbitrary, then
 $f(n) = n$, so f can reach any
element in \mathbb{N} .

- Bijective?

injective

Yes, as it is both ~~bijective~~
and surjective.

(b) If the domain / codomain of f are \mathbb{Z} ,
is f

- injective?

No, as $f(1) = f(-1) = 1$.

- Surjective?

No, as there is no $x \in \mathbb{Z}$ such
that $f(x) = -2$.

- Bijective?

No, as it is neither injective
nor surjective.

If the domain/codomain of f are \mathbb{R} , is
 f

- injective?

No, as again $f(1) = f(-1) = 1$.

- surjective?

Yes. Let $r \in \mathbb{R}$ be arbitrary. Then

- if $r \geq 1$, $f(r) = r$,

- if $0 \leq r < 1$, $f(\sqrt{r}) = r$, and

- if $r < 0$, $f\left(\frac{r-3}{2}\right) = r$,

so $\forall r$ we can find x such that $f(x) = r$.

- bijective?

No, as it is not injective.